

Relative Abundance, Size and Estimated Weight of Fish Collected at Hammond Lake

Common Name*	Number	Percent	Minimum	Maximum	Weight (lb)**	Percent
			Length (in)	Length (in)		
Golden shiner	83	31.0	3.4	8.5	9.62	8.8
Brown bullhead	50	18.7	2.4	14.8	20.88	19.0
Black crappie	48	17.9	2.2	10.7	2.77	2.5
Lake chubsucker	36	13.4	4.7	9.5	6.94	6.3
Bowfin	18	6.7	19.0	23.0	61.46	55.9
Bluegill	9	3.4	3.0	7.7	0.99	0.9
Grass pickerel	6	2.2	8.5	11.4	2.31	2.1
Largemouth bass	6	2.2	2.2	10.8	2.13	1.9
Pirate perch	4	1.5	2.6	3.4	0.02	0.0
Central mudminnow	2	0.7	3.4	3.7	0.02	0.0
Warmouth	2	0.7	6.5	7.5	0.55	0.5
Banded killifish	1	0.4	1.8	1.8	0.01	0.0
Redear	1	0.4	8.1	8.1	0.39	0.4
Spotted gar	1	0.4	22.0	22.0	1.39	1.3
Yellow bullhead	1	0.4	9.2	9.2	0.38	0.3
TOTAL	268				109.86	

*Common names of fishes recognized by the American Fisheries Society.

**Weights estimated from standard length-weight regression models.